

Scotland's Rural College

## Exposure of harbour seals (*Phoca vitulina*) to *Brucella* in declining populations across Scotland

Kershaw, JL; Stubberfield, EJ; Foster, G; Brownlow, A; Hall, AJ; Perrett, LL

*Published in:*  
Diseases of Aquatic Organisms

*DOI:*  
[10.3354/dao03163](https://doi.org/10.3354/dao03163)

First published: 20/09/2017

*Document Version*  
Peer reviewed version

[Link to publication](#)

### *Citation for pulished version (APA):*

Kershaw, JL., Stubberfield, EJ., Foster, G., Brownlow, A., Hall, AJ., & Perrett, LL. (2017). Exposure of harbour seals (*Phoca vitulina*) to *Brucella* in declining populations across Scotland. *Diseases of Aquatic Organisms*, 126(1), 13 - 23. <https://doi.org/10.3354/dao03163>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Exposure of harbour seals (*Phoca vitulina*) to *Brucella* in declining populations across Scotland**

\*Joanna L. Kershaw<sup>1</sup>, Emma J. Stubberfield<sup>2</sup>, Geoffrey Foster<sup>3</sup>, Andrew Brownlow<sup>3</sup>, Ailsa J. Hall<sup>1</sup>, Lorraine L. Perrett<sup>2</sup>.

1. Sea Mammal Research Unit, Scottish Oceans Institute, University of St Andrews, St Andrews, Fife, UK

\* jk49@st-andrews.ac.uk corresponding author

ajh@st-andrews.ac.uk

2. Animal and Plant Health Agency (APHA), Dept. of Bacteriology, New Haw, Addlestone, Surrey, UK

lorraine.perrett@apha.gsi.gov.uk

emma.stubberfield@apha.gsi.gov.uk

3. Scottish Marine Animals Strandings Scheme, SAC Consulting Veterinary Services, Inverness, UK

geoffrey.foster@sac.co.uk

andrew.brownlow@sac.co.uk

## ABSTRACT

Since 2000 there have been major declines in the abundance of Scottish harbour seals (*Phoca vitulina*). The causes of the declines remain uncertain. The aim of this study was to establish the extent to which the seals in the regions of greatest decline have been exposed to *Brucella*, a bacterial pathogen that causes reproductive failure in terrestrial mammalian hosts. Tissues from dead seals collected between 1992 and 2013 were cultured for *Brucella* (n=150). Serum samples collected from live capture-released seals (n=343) between 1997 and 2012 were tested for *Brucella* antibodies using the Rose Bengal plate agglutination test (RBT) and a competitive Enzyme-Linked Immunosorbent Assay (ELISA). 16% of seals cultured had *Brucella* isolated from one or more tissues but there were no pathological signs of infection. The cELISA results were more sensitive than the RBT results showing that overall, 25.4% of seals were seropositive with the highest seroprevalence in juveniles. As there was no evidence of either a higher seroprevalence, or higher circulating antibody levels in seropositive animals in the areas with the greatest declines, it was concluded that *Brucella* infection is likely not a major contributing factor to recent declines. However, the consistently high proportion of seals exposed to *Brucella* indicates possible endemicity in these populations, likely due to *Brucella pinnipedialis*, which has demonstrated a preference for pinniped hosts. Importantly, given the close proximity between seals, humans and livestock in many areas, there is the potential for cross-species infections.

## KEY WORDS

Pinnipeds, *Brucella*, disease, cultures, seroprevalence, antibodies, ELISA, Rose Bengal plate agglutination test

## INTRODUCTION

Aerial surveys have been carried out by the Sea Mammal Research Unit to monitor harbour seal (*Phoca vitulina*) populations around Scotland since 1985, and declines in a number of these populations have been seen since 2000 (Lonergan et al., 2007). Major declines of 68% in Orkney, 50% in Shetland, and 90% in the Firth of Tay have been documented in particular (SCOS, 2012). However, the pattern of the declines is not universal, as some areas remain more stable while the populations in other areas continue to decrease in size. The population in the Eden and Firth of Tay Special Area of Conservation, for example, has experienced the most dramatic and sustained declines of over 90% in the last 15 years with a most recent estimate of just 29 individuals left in 2014 (Hanson *et al.* 2015).

Many potential causes of the decline have been suggested, but the contributing factors remain uncertain. Some of these include predation by killer whales (Bolt et al., 2009), competition for food with other marine top predators (SCOS, 2012), exposure to biotoxins from harmful algal blooms (Hall and Frame, 2010), deliberate shooting (Thompson et al., 2007), accidental mortalities as a result of interactions with shipping vessels (Thompson et al., 2010), and predation by grey seals (*Halichoerus grypus*) (Brownlow *et al.* 2016). A further potential contributing factor to these declines is infectious disease, but there has been a lack of reports of sick animals by the Scottish Marine Animals Strandings Scheme (SMASS) or the Scottish Society for the Prevention of Cruelty to Animals (SSPCA). This suggests that if infectious disease was present in these populations, and was contributing to the observed declines, it could either be affecting the reproductive success of the animals, or, causing them to die very quickly once infected, or both. A particular infectious agent of interest in this respect is *Brucella* as it is known to cause reproductive failure in other mammalian hosts. The aim of this study was to establish the extent to which harbour seals in Scotland have been exposed to

*Brucella* over time, both before and during the observed population declines, and whether this could be a potential contributing factor to the major declines in some areas.

Members of the genus *Brucella* are Gram-negative, rod-shaped bacteria that cause chronic disease most commonly associated with abortions and infertility in domestic livestock (Seleem et al., 2010). Since the first reports of *Brucella* in a marine mammal in 1994 (Ross et al., 1994), infections have been recognised in a range of pinniped and cetacean species worldwide (Thakur et al., 2012). Strains isolated from marine mammals have been shown to be phenotypically and genetically distinct from those isolated from terrestrial mammals, and two species have been described that have pinnipeds and cetaceans as their preferred hosts, *Brucella pinnipedialis* and *Brucella ceti* respectively (Foster et al., 2007). *Brucella ceti* infections in cetaceans have been associated with various pathologies which include abortions and neonatal mortality (Miller et al., 1999), epididymitis in males (Dagleish et al., 2008), meningoencephalitis (González et al., 2002; Jauniaux et al., 2010; Alba et al., 2013; Garofolo et al., 2014), abscesses (Foster et al., 1996; Foster et al., 2002), endocarditis (González-Barrientos et al., 2010), mastitis, pneumonia, peritonitis, osteomyelitis and spinal discospondylitis (Foster et al., 2002). In contrast, pathology associated with *Brucella pinnipedialis* in seals is lacking despite several reports of its isolation (Foster et al., 2002; Nymo et al., 2011; Siebert et al., 2017; Tryland et al., 2005). With respect to harbour seals specifically, *Brucella* has previously been isolated from wild animals (Foster et al., 2002; Garner et al., 1997; Prenger-Berninghoff et al., 2008; Ross et al., 1994), and in several areas they have also been found to be seropositive (Gaydos et al., 2005; Hueffer et al., 2013; Maratea et al., 2003; Ross et al., 1996). However, the extent to which harbour seals in Scotland, or the United Kingdom in general, are currently exposed to this pathogen since the onset of the declines is unknown. Here, tissue samples from dead harbour seals collected by the SMASS over 20 years were cultured to test for *Brucella* isolates. In addition, harbour seal

serum samples from live-capture release studies by the Sea Mammal Research Unit over a 14 year period were tested for *Brucella* antibodies. Tempo-spatial patterns in seroprevalence were examined in order to investigate the potential role of *Brucella* as a contributing factor to the Scottish harbour seal declines.

## **MATERIALS & METHODS**

### ***Brucella* Cultures**

Tissue samples from 150 dead harbour seals from across Scotland, which had received post mortem examination were collected by the SMASS between 1992 and 2013 as part of systematic surveillance studies. Microbiological culture was performed including specific methods for *Brucella* isolation. The selected tissues varied between animals but typically included lung, liver, kidney, spleen, and small intestine, but also brain, pancreas, reproductive tissue, various lymph nodes and any abscesses apparent at post mortem. These were collected from approximately equal numbers of males (n = 61) and females (n = 41) and unsexed animals (n = 48). The majority of cases were adult animals although some juveniles were also sampled. Tissues were processed using a standardised method (Foster et al., 2002) and cultured on Columbia sheep blood agar (CSBA) (Oxoid, Basingstoke, UK) and Farrell's medium (Farrell, 1974) incubated at 37°C in air with 5% added CO<sub>2</sub>. Isolates with colonial appearance typical of *Brucella* on either medium were identified as *Brucella* using phenotypic tests as previously described (Foster et al., 2002). A multi-locus variable number of tandem repeats analysis (MLVA-16) was used to confirm species designation (Macquart et al., 2009).

### **Serum Sampling Procedure**

Blood samples were collected from 343 live-captured harbour seal adults, juveniles (<50kg and/or 120cm) and suckling pups from multiple haul out sites around five areas of Scotland; the South East (the Eden Estuary, the Firth of Tay and the Firth of Forth), the North East (the Moray Firth, Dornoch Firth, Loch Fleet and the Pentland Firth), the North West (the Isle of Skye, Loch Shiel, and the Loch Nan Uamh Islands), the South West (the Sound of Jura and south east Islay), and Orkney (Fig.1). Samples were collected between 1997 and 2012 at varying times of the year. Due to the opportunistic nature of the analysis of stored samples, sample sizes varied regionally, across years and between age classes although the male to female ratio was approximately equal with 182 males and 160 females (Table 1). The seals were captured in nets on haul outs or in the water, and were sedated with Zoletil 100 (Virbac, France) at a dose rate of 0.5 ml/100 kg body weight intravenously. Blood samples were taken from the extradural vein immediately after the immobilisation of the animal. The whole blood samples were spun, sera was collected and aliquots were frozen at -20°C for later analysis. Samples were collected under the Animal (Scientific Procedures) Act, 1986, Home Office Project and Personal Licences issued to the Sea Mammal Research Unit.

### **Serological Methods**

Rose Bengal plate agglutination test: In a preliminary trial, the Rose Bengal plate agglutination test (RBT) was used to test stored serum samples for the presence of *Brucella* antibodies. Serum samples were tested with the Micropath Rose Bengal kit against the *B. abortus* antigen (Omega Diagnostics, Alva, United Kingdom) following the kit instructions. Samples were either classed as positive or negative based on visually discernible agglutination of antigens. A positive and a negative control supplied by the kit were used for each set of four serum samples tested simultaneously.

Following the completion of the Rose Bengal trials with the successful detection of antibodies, the variation in seroprevalence, i.e. the proportion of animals with antibody levels higher than a background threshold level set for terrestrial mammals, as well as the absolute antibody levels in the samples, were investigated using a competitive ELISA.

Competitive ELISA (cELISA) - Polystyrene microtitre plates were coated with *Brucella melitensis* lipopolysaccharide (LPS) antigen. Both positive (positive serum from an experimentally infected goat with *B. melitensis*) and negative controls (no sera) were tested. Serum and a peroxidase labelled monoclonal antibody (BM40 from a locally held hybridoma) were added to the plates and incubated for 30 min at room temperature. The plate was then washed, and chromogen and substrate were added and incubated for a further 15 minutes at room temperature, shaking at 160rpm. The plates were read at an optical density (OD) of 450nm. The mean OD of duplicate wells was expressed as a percentage of antibodies binding to the plate. Test samples with an OD of less than 60% of the conjugate only control (no sera added) were recorded as positive. This cut-off threshold was established based on serology results from terrestrial mammals (Perrett et al., 2010). As such, a weak reaction, indicative of low antibody levels, was considered to be between 30-60% antibody binding, while a strong reaction, indicative of high circulating antibody levels, was considered to be <30%.

### **Statistical Analysis of Serological Data**

All statistical analyses were performed using the statistical package, R, version 3.1.2 (R Core Development Team, 2014). Statistical significance was taken at  $p = 0.05$ . Two different statistical approaches were taken to investigate firstly, variation in the seroprevalence data, and secondly, variation in the antibody levels in the seropositive individuals.

Seroprevalence data: Generalised linear models (glms) with a binomial distribution were fitted to the seroprevalence data with individuals classed as seropositive (1) and seronegative



(0) for the two tests separately. The 14 years of data was split into four time periods: 1997-2000 (n = 63) represents the years before the start of the decline, and the years between 2001 and 2012 were split into 3 periods with approximately equal numbers of samples in each to give the maximum statistical power for the analysis. These were 2001-2005 (n = 87), 2006-2008 (n = 94), 2009-2012 (n = 99). A global model including all explanatory variables of interest (region, sex, age-class, time period and an interaction between region and time period) was generated, and backwards variable selection using the 'step' function in the 'car' library in R (version 2.11.1) was performed to identify the combination of variables that best explained the variation in the data by producing the model with the lowest AIC.

Antibody levels data: In addition, variation in the levels of circulating *Brucella* antibodies were investigated in the seropositive individuals identified using the cELISA data. Antibody binding results of only the seropositive seals were modelled using a glm with a gamma distribution and a log-link function to model the non-normal distribution of the antibody binding data as most individuals had low circulating antibody levels while few were very high. Again, a global model with region, sex, age-class, time and an interaction between region and time was generated and backwards variable selection using the 'step' function was used to identify the combination of variables that best explained the variation in the data.

## RESULTS

### *Brucella* Cultures

Of the 150 animals examined bacteriologically between 1992 and 2013, *Brucella* was isolated from the tissues of 24 individuals (16 %). Details for 11 of these animals have been reported previously (Foster et al., 2002). None of the culture positive animals showed any signs of pathological lesions associated with infection and the cause of death was always associated with starvation, trauma or some other viral or bacterial infection, but not *Brucella*.

Of the tissues cultured, the two that were the most commonly culture positive were lung (45.8%) and spleen (41.7%) although not all tissues were sampled consistently across the individuals (Table 2). MLVA-16 analysis identified isolates as *B. pinnipedialis* belonging to one of two sequence types: ST 24 or ST 25. Serum was sampled from 12 of these culture positive seals, two of which were seronegative and were sampled from healthy animals that had been shot (Table 2).

## **Serology**

### **Test Performance**

The RBT trials were able to detect antibodies in the archived serum samples, and results showed that across all study sites over the whole sampling period, the prevalence of *Brucella* antibodies was 15.9%. However, the cELISA results showed a higher overall seroprevalence of 25.4%. The prevalence across all age and sex classes, as well as across regions and over time was lower for the RBT results compared to the cELISA results (Table 3).

All of the samples that were positive using the RBT were also classed as positive using the cELISA. These samples had the lowest antibody binding indicating the highest circulating *Brucella* antibodies. Specifically, the mean antibody binding of the samples classed as positive by the RBT was  $45.0 \pm 0.06\%$ , while those classed as negative had a mean antibody binding of  $74.0 \pm 0.02\%$  (Two sample t-test;  $t = 8.79$ ,  $df = 51.54$ ,  $p < 0.0001$ ). Therefore, it seems that the RBT is only able to detect high antibody levels in the serum samples, and is the least sensitive of the two serological methods tested. As a result, the RBT results may have underestimated the prevalence of *Brucella* antibodies in these harbour seals (Table 3).

### **Seroprevalence**

Using the cELISA seroprevalence data where individuals were classed as either seropositive or seronegative, the best binomial glm after backwards variable selection included only age class as an important explanatory variable. Juveniles had a significantly higher seroprevalence than adults and pups ( $p$  values  $< 0.025$ ), while adults and pups were not significantly different to each other ( $p = 0.11$ ). There were no significant changes in prevalence over time or between regions, and there were equal numbers of seropositive males and females.

### **Antibody Levels**

Variation in the levels of circulating *Brucella* antibodies were investigated in the seropositive individuals. Backwards variable selection of the glm using the cELISA antibody binding results of only the seropositive seals revealed that age class and an interaction between region and time period were retained in the final model with significant effects. Pups had near-significant higher antibody binding than both adults and juveniles ( $p$  values both  $< 0.07$ ), indicating the lowest circulating antibodies in these seropositive individuals (Fig. 2). There was no difference between the circulating antibody levels in juveniles and adults (Fig. 2). The interaction between region and time revealed that there were different patterns in circulating antibody levels in the seropositive seals between regions over the 14 year sampling period. The highest average circulating *Brucella* antibody levels (shown as the lowest % antibody binding in Fig. 3) were measured in the 2001-2005 time period in the South West and these then decreased over the following years ( $p = 0.035$ ). All areas showed a decrease followed by an increase again over the whole time frame with the exception of Orkney that showed a sustained decrease in circulating antibodies between 2001 and 2012 (Fig. 3), and the North West where seropositive individuals were only recorded in the final time period, but this is likely a reflection of very limited sampling before 2009 (Table 1) rather than a recent introduction of the bacteria to the area. Males and females had similar antibody levels.

## DISCUSSION

*Brucella* species were isolated from 16% of the dead stranded animals tested, but there were no signs of *Brucella*-specific pathological lesions associated with infection in these 24 animals. They all appear to have died of other causes, although it is possible that *Brucella* acted as a secondary infection in these cases. The results presented here therefore suggest that harbour seals can be infected by *Brucella*, likely *Brucella pinnipedialis*, without evidence of associated disease. *Brucella pinnipedialis* has also been cultured from apparently healthy tissues of a number of other pinniped species including grey seals (*Halichoerus grypus*) (Foster et al., 2002; Prenger-Berninghoff et al., 2008), hooded seals (*Cystophora cristata*) (Foster et al., 2002; Tryland et al., 2005), ringed seals (*Pusa hispida*) (Forbes et al., 2000) and harp seals (*Pagophilus groenlandicus*) (Forbes et al., 2000). At present, there is only limited evidence of *Brucella* infection causing disease in any species of phocid seal (Jauniaux et al., 2013), although *Brucella* isolation was suggested as a possible cause of abortion in an otariid species, the California sea lion (*Zalophus californianus*), with recovery of *Brucella* from the placenta and stomach contents of an aborted foetus (Goldstein et al., 2009). Extensive typing of a large number of marine mammal *Brucella* strains in a recent study, however, found the Californian Sea lion isolates to be similar to *B. ceti* recovered from bottlenose dolphins in the USA, and that *B. pinnipedialis* isolates from harbour seals in the USA were found to be closely related to Scottish strains (Whatmore et al., submitted for publication).

Animal experimentation in cattle with a *Brucella* isolate recovered from a Pacific harbour seal (*Phoca vitulina richardsii*) resulted in seroconversion and abortion in 2 of 3 pregnant animals, suggesting that some strains of *B. pinnipedialis* may have abortifacient potential (Rhyan et al., 2001). However, there has been no evidence in pinnipeds of disease due to *B. pinnipedialis* as has been seen with *B. ceti* infection in dolphins and porpoises where chronic

disease with significant clinical and pathological signs including male infertility, neurobrucellosis, cardiopathies, bone and skin lesions, and live strandings have been documented (Guzmán-Verri et al., 2012). While no evidence of disease has been reported in pinnipeds, detecting abortions in wild populations is very difficult, especially if the occurrence remains constant over time and there is limited data on pupping success, as is the case for the populations sampled here. The ability to detect abortions and monitor pupping success in different populations is therefore needed in order to determine that *B. pinnipedialis* does not cause disease in these seals.

Similar to this study, *Brucella* antibodies have been detected in sera from a number of marine mammal species using Rose Bengal tests (Hernández-Mora et al., 2008; Jensen et al., 2013; Retamal et al., 2000; Tryland et al., 1999; Tryland et al., 2005), and using both indirect ELISA (iELISA) and competitive ELISAs primarily designed for ruminants (Jensen et al., 2013; Lynch et al., 2011; Nielsen et al., 2001; Nymo et al., 2013a; Roe et al., 2010; Tachibana et al., 2006; Tryland et al., 1999; Van Bresseem et al., 2001). Here, differences in the prevalence estimates obtained from the RBT and the cELISA results highlight the need to consider test performance when conducting serological studies. It seems that the RBT is only able to detect antibodies when at higher levels, and as such, seropositive samples with low levels of antibody are not recognised.

Our results suggest that the cELISA appears to be a more sensitive test than RBT and is thus able to distinguish between seronegative samples and samples with low antibody levels. The cELISA results were therefore chosen for further analysis over the RBT results as this is thought to be a more robust and objective test. However, when detecting *Brucella* antibodies using serological methods, serological cross-reactions and false positives are potentially a major problem, and may contribute to the higher seroprevalence seen in the cELISA results. It is thought that in cattle, most problems caused by cross-reactivity are the result of

antibodies produced through the immune response of the animal to other microorganisms sharing similar structural characteristics with the O-polysaccharide of *Brucella* species (Corbel, 1985). We cannot rule out the possibility that other cross-reacting bacteria could affect these results. In cattle, it is thought that the cELISA is a more appropriate serological test than the iELISA as it is better able to distinguish between antibodies to *Brucella* species and antibodies to other cross-reacting Gram-negative bacteria (Nielsen, 1990; Samartino et al., 1999). In two studies on Australian fur seals (Lynch et al., 2011) and Hawaiian monk seals (*Monachus schauinslandi*) (Nielsen et al., 2005), it was concluded that the iELISA was an unreliable test for the identification of seropositive individuals. Thus, based on this previous evidence, the cELISA was chosen here as an appropriate assay as it is more conservative than an iELISA with a reduced chance of false positives.

The cELISA results indicate that approximately 25% of the seals sampled had antibodies to *Brucella*. This is within the range of previous studies on harbour seal populations in the North Atlantic where prevalence ranged between 3.1% (n=96) in the St Lawrence Estuary, 14% (n = 21) (Maratea et al. 2003) and 50% (n=8) off the Atlantic coast of the United States (Nielsen et al., 2001). Serology testing of 300 Scottish harbour seals prior to 2002 found 147 (49 %) to be positive (Foster et al. 2002). As there appears to have been no change in antibody prevalence over this 14 year sampling period, or between regions, these data suggest that *Brucella* may be endemic in Scottish harbour seals and exposure to the bacteria seems to have remained constant over the study period. For endemicity to occur, a high and regular rate of transmission of the bacteria is required within a population.

The transmission of *Brucella* in marine mammals is poorly understood as there is little evidence to support any particular route of infection in these species. It is likely that the routes of transmission are similar to those of terrestrial mammals, whereby transmission occurs through exposure to infected placenta, birth fluids and vaginal secretions as well as by

venereal spread (Young, 2006). As *Brucella* has been isolated from the reproductive organs of several cetacean species (Foster et al., 2002; González-Barrientos et al., 2010; Miller et al., 1999), and from an aborted foetus of a captive bottlenose dolphin (Ewalt et al., 1994), the most likely mode of transmission of *B. ceti* appears to be through sexual intercourse, vertical transmission from mother to foetus, maternal feeding and contact with aborted foetuses and placental tissues (Guzmán-Verri et al., 2012). The transmission between pinnipeds is even less well understood, but it could be similar to cetaceans. However, transmission may also occur through contact with infected individuals in gregarious species that haul out together in large groups. *Brucella* was cultured from the faeces of a seropositive juvenile harbour seal in captivity (Gaydos et al., 2005), suggesting that some *Brucella*-positive seals are actively shedding the bacteria. In addition, *B. pinnipedialis* was cultured or detected by PCR in harbour seal salivary gland secretions, lungs, urinary bladder, and faeces (Lambourn et al. 2013), suggesting that seals could be exposed to the bacterium via exposure to oral secretions, urine, or faeces on haul-outs. *Brucella* has also been isolated from subcutaneous lesions in cetaceans (Foster et al., 1996; Foster et al., 2002), so the potential for direct contact with similarly infected skin lesions, should they occur, in pinnipeds that haul out together may present another mode of transfer of the bacteria, although such lesions have not been reported to date. Together, this could make harbour seals more at risk of bacterial transfer at particular times during their life cycle when they haul-out in larger numbers during the breeding season and during the moult. Thus, the requirements of a high and regular rate of transmission of the bacteria for endemicity to occur could be met for harbour seals.

There was no regional variation in seroprevalence across Scotland, indicating that seals in the declining populations in Orkney and along the East coast have similar proportions of seropositive seals to the stable populations along the West coast. In addition there has been no change in the prevalence over time, even in declining populations, and none of the seals

sampled in this study showed any overt signs of ill health. These results further support the hypothesis that they may be infected by a strain of the bacteria that appears to be having little effect on their health. Other pinniped species have also been shown to be seropositive and yet remain apparently healthy and asymptomatic (Nielsen et al., 2005; Nielsen et al., 1996; Nymo et al., 2011; Retamal et al., 2000). Together, these results indicate that the bacteria may only cause a mild and transient infection, and *B. pinnipedialis* is most likely not a major cause of the harbour seal decline in Scotland. Other potential causes of the declines should therefore continue to be investigated.

Juveniles showed the highest overall prevalence of the three age classes. It has been reported that the higher incidence in juveniles may be as a result of recent exposure to the pathogen due to a change to a prey-based diet after they are weaned (Lynch et al., 2011; Nymo et al., 2013b). Lungworms carrying *Brucella* in fish prey species may be a means by which marine mammals become infected with the bacterium as was suggested when *Brucella* was isolated from the lungworms in a harbour porpoise (*Phocoena phocoena*) (Dawson et al., 2008). *Brucella* species have also been demonstrated by immunohistochemical staining in the uterus and the intestinal lumen of female *Parafulgaroides* lungworms from a Pacific harbour seal (Garner et al., 1997) and it was postulated that, based on the life cycle of the parasite, the larvae migrate through the respiratory tract and are then swallowed. From there, they pass through the digestive tract and out into the environment in the faeces where they are taken up by fish and ultimately by the seal. The parasitic larvae are released into the gastrointestinal tract of the animal, and when they mature into adults, they migrate to the lungs and continue the life cycle (Howard et al., 1983). It may also be significant that lung was the body tissue with the highest isolation rate from the 24 harbour seals that were positive by *Brucella* culture. As such, the high levels of antibodies in seropositive juveniles may suggest a more recent exposure to *Brucella* as they first start to eat fish containing the



infected parasites, but it is not necessarily indicative of an active infection. The seropositive adults have high antibody levels which may be indicative of both previous and regular exposure to the bacteria.

The finding that there were lower levels of *Brucella* antibodies in the seropositive pups compared to both seropositive adults and juveniles is surprising as it would be expected that passively transferred maternal antibodies would be present in pups, as they are found in the offspring of antibody-positive mothers in terrestrial species (Ray et al., 1988; Rhyan et al., 2009; Thakur et al., 2002). However, a lower seroprevalence was seen in Australian fur seal (Lynch et al., 2011), Hawaiian monk seal (Aguirre et al., 2007), hooded seal (Nymo et al., 2013b) and Alaskan harbour seal pups (Zarnke et al., 2006) compared to both adults and juveniles. In these studies, it was concluded that pups may have had maternal antibodies at titres lower than the threshold of detection used in their serological tests. These data support this theory that pups likely have low levels of maternal antibodies, and that they may not be exposed to infection until a later stage postweaning (Lynch et al., 2011). These findings further highlight the need for investigations into the timing of first exposure to *Brucella* and seroconversion as well as the development of specific thresholds of detection for antibodies to marine mammal strains of *Brucella* in various serological tests.

While the overall proportion of positive seals did not change across the different sampling regions over time, there were varying patterns of high and low antibody levels measured in the seropositive seals. Higher antibody levels were not recorded in the declining populations however, and there were no populations with consistently higher or lower antibody levels. The presence of antibodies does not necessarily suggest that the animals had a current or active infection at the time of sampling. The variation over time seen here between populations likely reflects cycles of infection followed by clearance in infected individuals that do not show any clinical signs of the disease. While the apparently high exposure rates of

Scottish harbour seals to *Brucella* appear not to be having a negative impact on their populations, such levels may have important implications for cross-species infections between humans and domestic livestock where infections may lead to disease. Currently a total of 53 marine mammal species worldwide have been shown to be seropositive for *Brucella* antibodies, and 20 of these species have been positive for *B. ceti* or *B. pinnipedialis* by culture or PCR assays (Foster et al., 2015; Hernández-Mora et al., 2013). The high seroprevalence seen here in all populations across Scotland suggests that wildlife professionals working with live seals could be exposed to the bacterium, and care should be taken when handling the animals and working with samples. To date, there have been four documented cases of humans infected with *Brucella ceti* (Brew et al., 1999; McDonald et al., 2006; Sohn et al., 2003), demonstrating the zoonotic potential of that species, but human infections with *B. pinnipedialis* have not been documented.

In conclusion, over a quarter of Scottish harbour seals have detectable levels of antibodies to *Brucella* which may indicate endemicity in these populations, possibly to a strain of the pathogen that has little effect on the health of individuals. These prevalence rates do not appear to explain the declines in Orkney and along the East coast as the prevalence in these areas is the same as in populations along the west coast that remain stable. The causes of the decline are likely to vary between regions and are probably due to a combination of factors, but *Brucella* infection does not appear to be one of them, based on our findings and comparison with seroprevalence rates for Scottish harbour seals before 2002 (Foster et al., 2002). Despite the routine use of the serological tests used here in many assessments of exposure to *Brucella* species, further validation of the tests for marine mammals is needed, and the discrepancies between the two test types here highlight the need for careful interpretation of the results.

## ACKNOWLEDGEMENTS

The authors would like to thank all the staff and students at the Sea Mammal Research Unit who were involved in the live capture-release fieldwork, with particular thanks to Simon Moss and Matthew Bivins. We also acknowledge the contributions of Harry Ross, Tony Patterson and Bob Reid for the post mortems of seals and provision of tissues for culture. Finally we would like to acknowledge the NERC National Capability Funding grant number SMRU 10001 for the funding.

## REFERENCES

- Aguirre AA, Keefe TJ, Reif JS, Kashinsky L, Yochem PK, Saliki JT, Stott JL, Goldstein T, Dubey JP, Braun R, Antonelis G (2007) Infectious Disease Monitoring of the Endangered Hawaiian Monk Seal. *J Wildl Dis* 43:229-241.
- Alba P, Terracciano G, Franco A, Lorenzetti S, Cocumelli C, Fichi G, Eleni C, Zygmunt MS, Cloeckaert A, Battisti A (2013). The presence of *Brucella ceti* ST26 in a striped dolphin (*Stenella coeruleoalba*) with meningoencephalitis from the Mediterranean Sea. *Vet Microbiol.* 31:158-63.
- Bolt HE, Harvey PV, Mandleberg L, Foote AD (2009) Occurrence of killer whales in Scottish inshore waters: temporal and spatial patterns relative to the distribution of declining harbour seal populations. *Aquat Conserv* 19: 671–675.
- Brew SD, Perrett LL, Stack JA, MacMillan AP, Staunton NJ (1999) Human exposure to *Brucella* recovered from a sea mammal. *Vet Rec* 144: 483.
- Brownlow A, Onoufriou J, Bishop A, Davison N, Thompson D (2016) Corkscrew Seals: Grey Seal (*Halichoerus grypus*) Infanticide and Cannibalism May Indicate the Cause of Spiral Lacerations in Seals. *PLoS ONE* 11(6): e0156464.

- 430 Corbel MJ (1985) Recent advances in the study of *Brucella* antigens and their serological  
431 cross-reactions. Veterinary Bulletin 55: 927–942.
- 432 Dagleish MP, Barley J, Finlayson J, Reid RJ, Foster G (2008) *Brucella ceti* Associated  
433 Pathology in the Testicle of a Harbour Porpoise (*Phocoena phocoena*). J Comp Pathol 139:  
434 54-59.
- 435 Dawson CE, Perrett LL, Stubberfield EJ, Stack JA, Farrelly SSJ, Cooley WA, Davison NJ,  
436 Quinney S (2008) Isolation and characterisation of *Brucella* from the lungworms of a harbour  
437 porpoise (*Phocoena phocoena*). J Wildl Dis 44: 237-246.
- 438 Ewalt DR, Payeur JB, Martin BM, Cummins DR, Miller WG (1994) Characteristics of a  
439 *Brucella* Species from a Bottlenose Dolphin (*Tursiops truncatus*). Journal of Veterinary  
440 Diagnostic Investigation 6: 448-452.
- 441 Forbes LB, Nielsen O, Measures L, Ewalt DR (2000) Brucellosis in Ringed Seals and Harp  
442 Seals from Canada. J Wildl Dis 36: 595-598.
- 443 Foster G, Jahans KL, Reid RJ, Ross HM (1996) Isolation of *Brucella* species from cetaceans,  
444 seals and an otter. Vet Rec 138: 583-586.
- 445 Foster G, MacMillan AP, Godfroid J, Howie F, Ross HM, Cloeckert A, Reid RJ, Brew S,  
446 Patterson IAP (2002) A review of *Brucella* sp. infection of sea mammals with particular  
447 emphasis on isolates from Scotland. Vet Microbiol 90: 563-580.
- 448 Foster G, Osterman BS, Godfroid J, Jacques I, Cloeckert A (2007) *Brucella ceti* sp. nov. and  
449 *Brucella pinnipedialis* sp. nov. for *Brucella* strains with cetaceans and seals as their preferred  
450 hosts. International Journal of Systematic and Evolutionary Microbiology 57: 2688-2693.

- 451 Foster G, Whatmore, AM, Dagleish MP, Baily JL, Deaville R, Davison NJ, Koylas MS,  
452 Perrett LL, Stubberfield EJ, Reid RJ, Brownlow AC (2015) Isolation of *Brucella ceti* from a  
453 long-finned pilot whale (*Globicephala melas*) and a Sowerby's beaked whale (*Mesoloden*  
454 *bidens*). J Wildl Dis 51: 868-871.
- 455 Garner MM, Lambourn DM, Jeffries SJ, Hall PB, Rhyan JC, Ewalt DR, Polzin LM, Cheville  
456 NF (1997) Evidence of *Brucella* infection in Parafilaroides lungworms in a Pacific harbor  
457 seal (*Phoca vitulina richardsi*). Journal of Veterinary Diagnostic Investigation 9: 298-303.
- 458 Garofolo G, Zilli K, Troiano P, Petrella A, Marotta F, Di Serafino G, Ancora M, Di  
459 Giannatale E (2014) *Brucella ceti* from two striped dolphins stranded on the Apulia  
460 coastline, Italy. J Med Microbiol. 63:325-9.
- 461 Gaydos JK, Norman SA, Lambourn D, Jeffries S, Raverty S, Leslie M, Lockwood S,  
462 DeGhetto D, Huckabee J, Ewalt D, Whaley J, Rowles T (2005) Should harbour seals with  
463 antibodies to *Brucella* be rehabilitated? Presentation at the 36th Annual Conference of the  
464 International Association of Aquatic Animal Medicine.
- 465 Godfroid J, Garin-Bastuji B, Saegerman C, Blasco JM (2013) Brucellosis in terrestrial  
466 wildlife. Rev Sci Tech 32: 27-42.
- 467 Godfroid J, Nielsen K, Saegerman C (2010) Diagnosis of Brucellosis in Livestock and  
468 Wildlife. Croat Med J 51: 296-305.
- 469 Goldstein, T, Zabka TS, Delong, RL, Wheeler EA, Ylitalo G, Bargu S, Silver M, Leighfield  
470 T, van Dolah F, Langlois G, Sidor I, Dunn JL, Gulland FMD (2009) The role of domoic acid  
471 in abortion and premature parturition of Californian sea lions (*Zalophus californianus*) on  
472 San Miguel Island. J Wildl Dis 45: 91-108.

- 473 González-Barrientos R, Morales JA, Hernández-Mora G, Barquero-Calvo E, Guzmán-Verri  
474 C, Chaves-Olarte E, Moreno E (2010) Pathology of Striped Dolphins (*Stenella coeruleoalba*)  
475 Infected with *Brucella ceti*. J Comp Pathol 142: 347-352.
- 476 González L, Patterson IA, Reid RJ, Foster G, Barberán M, Blasco JM, Kennedy S, Howie  
477 FE, Godfroid J, MacMillan AP, Schock A, Buxton D (2002) Chronic Meningoencephalitis  
478 Associated with *Brucella* sp. Infection in Live-stranded Striped Dolphins (*Stenella*  
479 *coeruleoalba*). J Comp Pathol 126: 147-152.
- 480 Guzmán-Verri C, González-Barrientos R, Hernández-Mora G, Morales JA, Baquero-Calvo E,  
481 Chaves-Olarte E, Moreno E (2012) *Brucella ceti* and Brucellosis in Cetaceans. Frontiers in  
482 Cellular and Infection Microbiology 2: 1-22.
- 483 Hall AJ, Frame E (2010) Evidence of domoic acid exposure in harbour seals from Scotland:  
484 A potential factor in the decline in abundance? Harmful Algae 9: 489-493.
- 485 Hanson N, Thompson D, Duck C, Baxter J, Lonergan M (2015) Harbour seal (*Phoca*  
486 *vitulina*) abundance within the Firth of Tay and Eden estuary, Scotland: recent trends and  
487 extrapolation to extinction. Aquat Conserv Mar Freshw Ecosyst. DOI: 10.1002/aqc.2609
- 488 Hernández-Mora G, González-Barrientos R, Moreno E (2008) Neurobrucellosis in Stranded  
489 Dolphins, Costa Rica. Emerg Infect Dis 14: 1825.
- 490 Hernández-Mora G, Palacios-Alfaro JD, González-Barrientos R (2013) Wildlife reservoirs of  
491 brucellosis: *Brucella* in aquatic environments. Rev sci tech 32 : 89-103.
- 492 Howard EB, Britt JO, Matsumoto G (1983) Parasitic Diseases. Pathobiology of marine  
493 mammal diseases. (E.B. Howard, ed.). CRC Press, Boca Raton, Florida, 128–213.

- 494 Hueffer K, Gende SM, O'Hara TM (2013) Assay dependence of *Brucella* antibody  
495 prevalence in a declining Alaskan harbor seal (*Phoca vitulina*) population. Acta Vet Scand  
496 55: 1-5.
- 497 Jauniaux T, Brenez C, Frentin D, Godfroid J, Haelters J, Jacques T, Kerckhof F, Mast J,  
498 Sarlet M, Coignoul F (2010) *Brucella ceti* infection in harbor porpoise (*Phocoena*  
499 *phocoena*). Emerg Infect Dis 16: 1966–1968.
- 500 Jauniaux T, Didier M, Fretin D, Godfroid J, Laurent E, Michel P, Sarlet M, Van Gompel J,  
501 Coignoul F (2013) Brucellosis in two seal pups. 27th annual conference of the European  
502 Cetacean Society.
- 503 Jensen SK, Nymo IH, Forcada J, Hall AJ, Godfroid J (2013) *Brucella* antibody  
504 seroprevalence in Antarctic seals (*Arctocephalus gazella*, *Leptonychotes weddellii* and  
505 *Mirounga leonina*). Dis Aquat Org 105: 175-181.
- 506 Lambourn DM, Garner M, Ewalt D, Raverty S, Sidor I, Jeffries SJ, Rhyan J, Gaydos JK  
507 (2013) *Brucella pinnipedialis* infections in Pacific harbor seals (*Phoca vitulina richardsi*)  
508 from Washington State, USA. J Wildl Dis 49 : 802-815.
- 509 Lonergan M, Duck CD, Thompson D, Mackey BL, Cunningham L, Boyd IL (2007) Using  
510 sparse survey data to investigate the declining abundance of British harbour seals. J Zool  
511 271: 261-269.
- 512 Lynch M, Duignan PJ, Taylor T, Nielsen O, Kirkwood R, Gibbens J, Arnould JPY (2011)  
513 Epizootiology of *Brucella* Infection in Australian Fur Seals. J Wildl Dis 47: 352-363.
- 514 Macquart M, Le Fleche P, Foster G, Tryland M, Ramisse F, Djonne B, Al Dahouk S, Jacques  
515 I, Neubauer H, Walravens K, Godfroid J, Cloeckaert A, Vergnaud G (2009) MLVA-16  
516 typing of 295 marine mammal *Brucella* isolates from different animal and geographic origins

- 517 identifies 7 major groups within *Brucella ceti* and *Brucella pinnipedialis*. BMC Microbiol 9:  
518 145.
- 519 Maratea J, Ewalt D, Frasca S, Dunn JL, De Guise S, Szkudlarek L, St Aubin DJ, French RA  
520 (2003) Evidence of *Brucella* sp. Infection in Marine Mammals Stranded along the Coast of  
521 Southern New England. J Zoo Wildl Med 34: 256-261.
- 522 McDonald WL, Jamaludin R, Mackereth G, Hansen M, Humphrey S, Short P, Taylor T,  
523 Swinger J, Dawson CE, Whatmore AM, Stubberfield E, Perrett LL, Simmons G (2006)  
524 Characterization of a *Brucella* sp. Strain as a Marine-Mammal Type despite Isolation from a  
525 Patient with Spinal Osteomyelitis in New Zealand. J Clin Microbiol 44: 4363-4370.
- 526 Miller WG, Adams LG, Ficht TA, Cheville NF, Payeur JP, Harley DR, House C, Ridgway  
527 SH (1999) *Brucella*-Induced Abortions and Infection in Bottlenose Dolphins (*Tursiops*  
528 *truncatus*). J Zoo Wildl Med 30: 100-110.
- 529 Moreno E, Moriyón I (2006) The genus *Brucella* in the Prokaryotes. . Vol. V. (M Dworkin, S  
530 Falkow, E Rosenberg, KH Schleifer & E Stackebrandt, eds). Springer-Verlag, New York,  
531 315–456.
- 532 Nielsen K (1990) The Serological Response of Cattle Immunized with *Yersinia enterocolitica*  
533 O:9 or O:16 to *Yersinia* and *Brucella abortus* Antigens in Enzyme Immunoassays. Vet  
534 Immunol Immunopathol 24 : 373-382.
- 535 Nielsen O, Nielsen K, Braun R, Kelly L (2005) A comparison of four serologic assays in  
536 screening for *Brucella* exposure in Hawaiian monk seals. J Wildl Dis 41: 126-133.
- 537 Nielsen O, Nielsen K, Stewart REA (1996) Serologic Evidence of *Brucella* spp. Exposure in  
538 Atlantic Walruses (*Odobenus rosmarus rosmarus*) and Ringed Seals (*Phoca hispida*) of  
539 Arctic Canada. Arctic 49: 383-386.



- 540 Nielsen O, Stewart REA, Nielsen K, Measures L, Duignan PJ (2001) Serological Survey of  
541 *Brucella* spp. Antibodies in Some Marine Mammals of North America J Wildl Dis 17: 89-  
542 100.
- 543 Nymo IH, Godfroid J, Åsbakk K, Larsen AK, das Neves CG, Rødven R, Tryland M (2013a)  
544 A protein A/G indirect enzyme-linked immunosorbent assay for the detection of anti-  
545 *Brucella* antibodies in Arctic wildlife. Journal of Veterinary Diagnostic Investigation 25:  
546 369-375.
- 547 Nymo IH, Tryland M, Frie AK, Haug T, Foster G, Rodven R, Godfroid J (2013b) Age-  
548 dependent prevalence of anti-*Brucella* antibodies in hooded seals *Cystophora cristata*. Dis  
549 Aquat Org 106: 187-196.
- 550 Nymo IH, Tryland M, Godfroid J (2011) A review of *Brucella* infection in marine mammals,  
551 with special emphasis on *Brucella pinnipedialis* in the hooded seal (*Cystophora cristata*). Vet  
552 Res 42.
- 553 Perrett LL, McGiven JA, Brew S, Stack JA (2010) Evaluation of Competitive ELISA for  
554 Detection of *Brucella* Infection in Domestic Animals. Croatia Medical Journal 51: 316-321.
- 555 Prenger-Berninghoff E, Siebert U, Stede M, Koenig A, Weiss R, Baljer G (2008) Incidence  
556 of *Brucella* species in marine mammals of the German north sea. Dis Aquat Org 81: 65-71.
- 557 Ray WC, Brown RR, Stringfellow DA, Schnurrenberger PR, Scanlan CM, Swann AI (1988)  
558 Bovine brucellosis: an investigation of latency in progeny of culture-positive cows. J Am Vet  
559 Med Assoc 15: 182-186.
- 560 Retamal P, Abalos P, Blank O, Torres D (2000) Detection of anti- *Brucella* antibodies in  
561 pinnipeds from the Antarctic territory. Vet Rec 146: 166-167.

- 562 Rhyan JC, Aune K, Roffe T, Ewalt D, Hennager S, Gidlewski T, Olsen S, Clarke R (2009)  
563 Pathogenesis and epidemiology of Brucellosis in Yellowstone Bison: Serologic and culture  
564 results from females and their progeny. J Wildl Dis 45: 729-739.
- 565 Roe WD, Rogers LE, Gartrell BD, Chilvers BL, Duignan PJ (2010) Serologic Evaluation of  
566 New Zealand Sea Lions for Exposure to *Brucella* and *Leptospira* spp. J Wildl Dis 46: 1295-  
567 1299.
- 568 Ross HM, Foster G, Reid RJ, Jahans KL, MacMillan AP (1994) *Brucella* species infection in  
569 sea-mammals. Vet Rec 134: 359.
- 570 Ross HM, Jahans KL, MacMillan AP, Reid RJ, Thompson PM, Foster G, (1996) *Brucella*  
571 species infection in North Sea seal and cetacean populations. Vet Rec 138: 647-  
572 648. Saegerman C, Berkven D, Godfroid J, Walravens K (2010) Bovine brucellosis. In:  
573 Infectious and parasitic diseases of livestock. Lavoisier, Paris.
- 574 Samartino L, Gall D, Gregoret R, Nielsen K (1999) Validation of enzyme-linked  
575 immunosorbent assays for the diagnosis of bovine brucellosis. Vet Microbio 70: 193-200.
- 576 SCOS (2012) Scientific Advice on Matters Related to the Management of Seal Populations:  
577 2012. SCOS Main Advice 2012.
- 578 Seleem MN, Boyle SM, Sriranganathan N (2010) Brucellosis: A re-emerging zoonosis. Vet  
579 Microbio 140: 392-398.
- 580 Siebert U, Rademaker M, Ulrich SA, Wohlsein P, Ronnenberg K, Prenger-Berninghoff E  
581 (2017) Bacterial microbiota in harbour seals (*Phoca vitulina*) from the North Sea of  
582 Schleswig-Holstein, Germany, around the time of morbillivirus and influenza epidemics. J  
583 Wildl Dis In-press

- 584 Sohn AH, Probert WS, Glaser CA, Gupta N, Bollen AW, Wong JD, Grace EM, McDonald  
 585 WC (2003) Human Neurobrucellosis with Intracerebral Granuloma Caused by a Marine  
 586 Mammal *Brucella* spp. Emerg Infect Dis 9.
- 587 Tachibana M, Watanabe K, Kim S, Omata Y, Murata K, Hammond T, Watarai M (2006)  
 588 Antibodies to *Brucella* spp. in Pacific Bottlenose Dolphins from the Solomon Islands. J Wildl  
 589 Dis 42: 412-414.
- 590 Thakur SD, Kumar R, Thapliyal DC (2002) Human brucellosis: review of an under-  
 591 diagnosed animal transmitted disease. J Commun Dis 34 : 287-301.
- 592 Thakur SD, Vaid RK, Panda AK, Saini Y (2012) Marine mammal brucellosis: a new  
 593 dimension to an old zoonosis. Curr Sci 103: 902-910.
- 594 Thompson D, Bexton S, Brownlow A, Wood D, Patterson T, Pye K, Lonergan M, Milne R  
 595 (2010) Report on recent seal mortalities in UK waters caused by extensive lacerations. Sea  
 596 Mammal Research Unit Reports.
- 597 Thompson PM, Mackey B, Barton TR, Duck C, Butler JRA (2007) Assessing the potential  
 598 impact of salmon fisheries management on the conservation status of harbour seals (*Phoca*  
 599 *vitulina*) in north-east Scotland. Anim Conserv 10: 48-56.
- 600 Tryland M, Kleivane L, Alfredsson A, Kjeld M, Arnason A, Stuen S, Godfroid J (1999)  
 601 Evidence of *Brucella* infection in marine mammals in the North Atlantic Ocean. Vet Rec  
 602 144: 588-592.
- 603 Tryland M, Sørensen KK, Godfroid J (2005) Prevalence of *Brucella pinnipediae* in healthy  
 604 hooded seals (*Cystophora cristata*) from the North Atlantic Ocean and ringed seals (*Phoca*  
 605 *hispida*) from Svalbard. Vet Microbio 105: 103-111.

606 Van BresseM M-F, Van Waerebeek K, Raga JA, Godfroid J, Brew SD, MacMillan AP  
607 (2001) Serological evidence of *Brucella* species infection in odontocetes from the south  
608 Pacific and the Mediterranean. Vet Rec 148: 657-661.

609 Whatmore AM, Dawson C, Muchowski J, Perrett LL, Stubberfield E, Koylass M, Foster G,  
610 Davison NJ, Quance C, Sidor IF, Field CL, St. Leger J Phenotypic and Molecular  
611 Characterisation of North American *Brucella* from Marine Mammals: Novel Genotypes, New  
612 Host Species Associations and Diagnostic Implications. Submitted for publication.

613 Young EJ (2006) *Brucella* spp., In: Gillespie SH, Hawkey PM (eds.) Principles and Practice  
614 of Clinical Bacteriology, Second Edition. John Wiley & Sons, Chichester.

615 Zarnke RL, Saliki JT, Macmillan AP, Brew SD, Dawson CE, Ver Hoef JM, Frost KJ, Small  
616 RJ (2006) Serologic survey for *Brucella* spp., phocid herpesvirus-1, phocid herpesvirus-2,  
617 and phocine distemper virus in harbour seals from Alaska, 1976-1999. J Wildl Dis 42: 290-  
618 300.

619

620

621


622

623


624

625


626



**Fig 1. Sampling regions of live-captured harbour seals across Scotland between 1997 and 2012 by the Sea Mammal Research Unit. Serum samples were grouped into 5 main regions across Scotland as indicated by the circles on the map. Over this time period, the populations along the west coast, marked as 'North West' and 'South West' were stable while the populations in 'Orkney', the 'North East' and the 'South East' underwent precipitous declines (SCOS, 2012).**



**Fig. 2. cELISA antibody binding of the seropositive animals by age class.** Low antibody binding indicates high levels of circulating *Brucella* antibodies in the seals. The pups had lower circulating antibody levels than both adults and juveniles.



**Fig. 3. Interaction plot of the cELISA % antibody binding over time for each sampling region showing variation in the seropositive individuals. Low antibody binding indicates high levels of circulating *Brucella* antibodies in the seals. The North West sampling region is not included here as seropositive animals were only identified between 2009 – 2012. With the exception of Orkney, the other sampling regions showed a decrease in the circulating levels of *Brucella* antibodies, followed by an increase again over this time period.**

**Table 1** – Serological samples collected from harbour seals across Scotland across the five sampling regions and over 12 years broken down into 4 time periods. Samples are grouped by sex and age class. A total of 306 adults, 15 juveniles and 22 pups were sampled.

Sampling Period	Sex	Age Class	Serological Samples Collected by Region				
			North East	North West	Orkney	South East	South West
1997 - 2000	Males	Adults	-	-	-	37	-
		Juveniles	-	-	-	1	-
		Pups	-	-	-	3	-
	Females	Adults	-	-	-	21	-
		Juveniles	-	-	-	1	-
		Pups	-	-	-	-	-
2001 - 2005	Males	Adults	5	-	11	15	14
		Juveniles	-	-	-	1	-
		Pups	-	-	-	-	-
	Females	Adults	10	-	12	9	4
		Juveniles	1	-	-	2	2
		Pups	-	-	-	-	1
2006 - 2008	Males	Adults	3	1	10	16	8
		Juveniles	-	-	-	-	1
		Pups	-	-	-	-	5
	Females	Adults	7	1	15	6	7
		Juveniles	1	-	-	-	-
		Pups	2	-	11	-	-
2009 - 2012	Males	Adults	1	14	21	10	2
		Juveniles	1	-	1	-	1
		Pups	-	-	-	-	-
	Females	Adults	6	11	20	1	8
		Juveniles	1	-	-	-	1
		Pups	-	-	-	-	-



**Table 2** - Details of 24 *Brucella* culture positive harbour seals. The last two digits of the reference number indicate the year of stranding. \* Indicates individuals that were culture positive but seronegative. MLN, mesenteric lymph node; IILN, internal iliac lymph node; EILN, external iliac lymph node; GLN, gastric lymph node, ManLN, mandibular lymph node; HLN, hepatic lymph node; TLN, thoracic lymph node and CLRN, colorectal lymph node; SI, small intestine.

Reference Number	Positive Cultures	Negative Cultures
M2357/93	Spleen	lung, liver
M2466/93	Spleen	lung, MLN
M2533/93	Spleen	MLN, SI
M292/94*	Spleen	testes, MLN, SI
M336/94*	IILN	spleen, MLN
M339/94	GLN	spleen, IILN
M972/94	EILN, manLN	spleen, MLN
M490/95	EILN, HLN, IILN, TLN	lung, spleen, brain, CRLN, GLN, manLN, blood, SI
M514/96	Lung	
M445/99	Lung	liver, spleen, kidney, brain, MLN, blood, SI
M13/01	Lung	liver, spleen, kidney, blood
M250/02	lung, liver, spleen, kidney, MLN, blood	brain, SI
M305/02	Spleen	lung, liver, kidney, brain, MLN, cellulitis, SI
M342/02	MLN	lung, liver, spleen, kidney, brain, blood
M374/02	lung, liver, spleen, kidney	brain, MLN
M449/02	Lung	
M599/02	MLN	lung, liver, spleen, kidney
M43/09	lung, liver, spleen, kidney, MLN, SI	
M91/10	lung, brain	liver, spleen, kidney, MLN
M228/10	pancreas	lung, liver, spleen, kidney, brain,

Scottish harbour seals' exposure to *Brucella*

		MLN, SI
M244/10	lung, liver, spleen, brain, MLN, SI	Kidney
M273/10	lung, MLN	liver, spleen, kidney, brain, pre- scapular LN, SI
M341/11	lung, spleen, MLN, SI	liver, kidney, brain, abscess
M337/13	kidney, brain	lung, liver, spleen

661

662

663

664

665

666

667

668

669

670

671

672

673

674

675

**Table 3** - Comparison of the prevalence (% of seropositive animals) using the Rose Bengal plate agglutination test (RBT) and the competitive ELISA (cELISA). The ELISA results indicate a higher overall prevalence of *Brucella* antibodies in harbour seals than the Rose Bengal results.

	% of Seropositive Animals	
	RBT	cELISA
<b>Scotland</b> (n = 343)	15.9	25.4
<b>North East</b> (n = 38)	28.6	28.9
<b>North West</b> (n = 27)	14.8	33.3
<b>Orkney</b> (n = 101)	12.9	28.7
<b>South East</b> (n = 123)	19.8	20.3
<b>South West</b> (n = 54)	3.9	24.1
<b>Males</b> (n = 183)	14.5	25.7
<b>Females</b> (n = 160)	17.4	24.4
<b>Adults</b> (n = 306)	14.5	25.2
<b>Juveniles</b> (n = 15)	60.0	53.3
<b>Pups</b> (n = 22)	4.5	9.09
<b>1997-2000</b> (n = 63)	12.3	22.2
<b>2001-2005</b> (n = 87)	24.7	26.4
<b>2006-2008</b> (n = 94)	12.0	18.1
<b>2009-2012</b> (n = 99)	13.7	33.3